

DUAL -1 HAHN POLYNOMIALS: "CLASSICAL" POLYNOMIALS BEYOND THE LEONARD DUALITY

SATOSHI TSUJIMOTO, LUC VINET, AND ALEXEI ZHEDANOV

ABSTRACT. We introduce the -1 dual Hahn polynomials through an appropriate $q \rightarrow -1$ limit of the dual q -Hahn polynomials. These polynomials are orthogonal on a finite set of discrete points on the real axis, but in contrast to the classical orthogonal polynomials of the Askey scheme, the -1 dual Hahn polynomials do not exhibit the Leonard duality property. Instead, these polynomials satisfy a 4-th order difference eigenvalue equation and thus possess a bispectrality property. The corresponding generalized Leonard pair consists of two matrices A, B each of size $N + 1 \times N + 1$. In the eigenbasis where the matrix A is diagonal, the matrix B is 3-diagonal; but in the eigenbasis where the matrix B is diagonal, the matrix A is 5-diagonal.

1. INTRODUCTION

Recently new explicit families of "classical" orthogonal polynomials $P_n(x)$ were introduced [12], [13], [14]. These polynomials satisfy an eigenvalue equation of the form

$$(1.1) \quad LP_n(x) = \lambda_n P_n(x).$$

The operator L is of first order in the derivative operator ∂_x and contains moreover the reflection operator R defined by $Rf(x) = f(-x)$; it can be identified as a first order operator of Dunkl type written as

$$(1.2) \quad L = F(x)(I - R) + G(x)\partial_x R$$

with some real rational functions $F(x), G(x)$. The corresponding polynomial eigensolutions $P_n(x)$ can be obtained from the big and little q -Jacobi polynomials by an appropriate limit $q \rightarrow -1$.

In [11] we generalized this approach to the case of Dunkl shift operators. In this case the operator L contains the shift operator $T^+f(x) = f(x + 1)$ together with the reflection operator R :

$$(1.3) \quad L = F(x)(I - R) + G(x)(T^+R - I)$$

(I stands for the identity operator). The rational functions can be recovered from the condition that the operator L stabilizes the spaces of polynomials, i.e. it sends any polynomial of degree n into a polynomial of the same degree. It can then be demonstrated [11] that the polynomial eigensolutions $P_n(x)$ of the eigenvalue equation (1.1) with the operator (1.3) satisfy the 3-term recurrence relation and hence are orthogonal polynomials. In fact, the polynomials $P_n(x)$ in this case coincide with the Bannai-Ito (BI) polynomials first constructed in [1] (see also [10]).

Key words and phrases. Classical orthogonal polynomials, dual q -Hahn polynomials, Leonard duality. AMS classification: 33C45.

The BI polynomials thus possess the bispectrality property: they satisfy simultaneously the 3-term recurrence relation (common to all orthogonal polynomials)

$$(1.4) \quad P_{n+1}(x) + b_n P_n(x) + u_n P_{n-1}(x) = x P_n(x)$$

and the eigenvalue equation (1.1). Moreover, in the case when the support of the orthogonality measure consists of a finite number of points x_s , $s = 0, 1, 2, \dots, N$, the BI polynomials satisfy the Leonard duality property [6], [1], [10]. This means that there is a finite difference equation of the form

$$(1.5) \quad U_s (P_n(x_{s+1}) - P_n(x_s)) + V_s (P_n(x_{s-1}) - P_n(x_s)) = \lambda_n P_n(x_s)$$

with some real coefficients V_s, U_s . The difference equation (1.5) is of the second order and can be considered as a dual relation with respect to the recurrence relation. In fact, the difference equation (1.5) is a simple consequence of the eigenvalue equation (1.1) with the Dunkl shift operator (1.3) [11].

We showed in [11] that the BI polynomials can be obtained by an appropriate $q \rightarrow -1$ limit from the Askey-Wilson polynomials. Correspondingly, the Dunkl shift operator L appears in the same limit from the Askey-Wilson difference operator [11]. It should be stressed that there are several possibilities in taking the limit $q \rightarrow -1$ of the Askey-Wilson polynomials. Not all of them lead to the BI polynomials. There is another family of orthogonal polynomials - called the complementary BI polynomials in [11] - which can be obtained from the Askey-Wilson polynomials in the same limit. In contrast to the BI polynomials, the complementary BI (CBI) polynomials do not satisfy an eigenvalue equation of the form (1.1). As a consequence, the CBI polynomials do not possess the Leonard duality property (1.5).

The main purpose of the present paper is to study the orthogonal polynomials which appear in the $q \rightarrow -1$ limit of the dual q -Hahn polynomials. We call these polynomials the dual -1 Hahn polynomials. We derive explicitly basic properties and relations for them including a 3-term recurrence relation. The main result is the existence of a dual eigenvalue equation (1.1) for the dual -1 Hahn polynomials. In contrast to the BI polynomials, the operator L is now 2-nd order with respect to the shift operators. As a consequence, the dual -1 Hahn polynomials obey a 5-term difference relation on an appropriate grid x_s (instead of the 3-term relation (1.5) for the BI polynomials).

2. DUAL q -HAHN POLYNOMIALS

The dual q -Hahn polynomials [5]

$$(2.1) \quad R_n(x; a, b, N) = {}_3\Phi_2 \left(\begin{matrix} q^{-n}, q^{-s}, abq^{s+1} \\ aq, q^{-N} \end{matrix} \middle| q; q \right)$$

depend on 3 parameters a, b, N , where N is a positive integer. The argument x can be parametrized in terms of s as follows

$$(2.2) \quad x = q^{-s} + abq^{s+1}.$$

The polynomials $R_n(x; a, b, N)$ satisfy the 3-term recurrence relation

$$(2.3) \quad A_n R_{n+1}(x; a, b, N) - (A_n + C_n - 1 - abq) R_n(x; a, b, N) + C_n R_{n-1}(x; a, b, N) = x R_n(x; a, b, N),$$

where

$$(2.4) \quad A_n = (1 - q^{n-N})(1 - aq^{n+1}), \quad C_n = aq(1 - q^n)(b - q^{n-N-1}).$$

The polynomials $R_n(x; a, b, N)$ are not monic; their monic version $P_n(x; a, b, N) = \kappa_n R_n(x; a, b, N) = x^n + O(x^{n-1})$ (with κ_n an appropriate factor) will obey the recurrence relation

$$(2.5) \quad P_{n+1}(x; a, b, N) + b_n P_n(x; a, b, N) + u_n P_{n-1}(x; a, b, N) = x P_n(x; a, b, N),$$

where

$$b_n = 1 + abq - A_n - C_n, \quad u_n = A_{n-1} C_n.$$

The dual q-Hahn polynomials also verify a q-difference equation of second order [5]

$$(2.6) \quad B(s)R_n(x_{s+1}) + D(s)R_n(x_{s-1}) - (B(s) + D(s))R_n(x_s) = (q^{-n} - 1)R_n(x_s), \quad s = 0, 1, 2, \dots, N,$$

where x_s is given by (2.2) and

$$(2.7) \quad \begin{aligned} B(s) &= \frac{(1 - q^{s-N})(1 - aq^{s+1})(1 - abq^{s+1})}{(1 - abq^{2s+1})(1 - abq^{2s+2})} \\ D(s) &= -\frac{aq^{s-N}(1 - q^s)(1 - abq^{s+1+N})(1 - bq^s)}{(1 - abq^{2s+1})(1 - abq^{2s})}. \end{aligned}$$

As per [7], the equation (2.6) can be interpreted as a q-difference equation on the "q-quadratic grid" x_s .

The dual q-Hahn polynomials satisfy the orthogonality relation

$$(2.8) \quad \sum_{s=0}^N w_s R_n(x_s) R_m(x_s) = \kappa_0 u_1 u_1 \dots u_n \delta_{nm},$$

where the discrete weights are

$$(2.9) \quad w_s = \frac{(aq, abq, q^{-N}; q)_s}{(q, abq^{N+2}, bq; q)_s} \frac{1 - abq^{2s+1}}{(1 - abq)(-aq)^s} q^{Ns - s(s-1)/2}$$

and the normalization constant is

$$\kappa_0 = \frac{(abq^2; q)_N}{(bq; q)_N} (aq)^{-N}.$$

We used the standard notation for the q-shifted factorials [5].

When $q \rightarrow 1$ and $a = q^\alpha$, $b = q^\beta$ the dual q-Hahn polynomials become the ordinary dual Hahn polynomials [5]

$$(2.10) \quad W_n(x_s; a, b, N) = {}_3F_2 \left(\begin{matrix} -n, -s, s+1+\alpha+\beta \\ \alpha+1, -N \end{matrix} \middle| 1 \right)$$

where

$$x_s = s(s + \alpha + \beta + 1).$$

These polynomials satisfy the three-term recurrence relation

$$(2.11) \quad A_n W_{n+1}(x_s) - (A_n + C_n) W_n(x_s) + C_n W_{n-1}(x_s) = x_s W_n(x_s),$$

where

$$A_n = (n - N)(n + \alpha + 1), \quad C_n = n(n - \beta - N - 1).$$

The corresponding monic dual Hahn polynomials $\hat{W}_n(x)$ obey

$$(2.12) \quad \hat{W}_{n+1}(x_s) + b_n \hat{W}_n(x_s) + u_n \hat{W}_{n-1}(x_s) = x_s \hat{W}_n(x_s),$$

where

$$(2.13) \quad u_n = A_{n-1} C_n = n(n - \beta - N - 1)(n - N - 1)(n + \alpha), \quad b_n = -A_n - C_n.$$

3. A LIMIT $q \rightarrow -1$ OF THE RECURRENCE RELATIONS AND THE ORTHOGONALITY PROPERTY

We wish to consider a limit $q \rightarrow -1$. We will assume that $a \rightarrow \pm 1$ and $b \rightarrow \pm 1$ when $q \rightarrow -1$. We want to obtain a nondegenerate limit of the coefficients $A_n(1+q)^{-1}, C_n(1+q)^{-1}$ for $q \rightarrow -1$. This means that both these limit coefficients should exist and be nonzero for all admissible values $n = 1, 2, \dots, N$. It is hence easily seen that necessarily, we must have $ab \rightarrow 1$. Two situations have to be considered separately:

(i) when $N = 2, 4, 6, \dots$ is even, the nontrivial $q \rightarrow -1$ limit then exists iff $a \rightarrow 1, b \rightarrow 1$. It is then natural to take the parametrization

$$(3.1) \quad q = -e^\varepsilon, \quad a = e^{-\alpha\varepsilon}, \quad b = e^{-\beta\varepsilon}, \quad \varepsilon \rightarrow 0$$

with real parameters α, β . Dividing the recurrence relation (2.3) by $q+1$ and taking the limit $\varepsilon \rightarrow 0$ we obtain the recurrence relation

$$(3.2) \quad A_n^{(-1)} R_{n+1}^{(-1)}(y_s; \alpha, \beta, N) + C_n^{(-1)} R_{n-1}^{(-1)}(y_s; \alpha, \beta, N) - (A_n^{(-1)} + C_n^{(-1)}) R_n^{(-1)}(y_s; \alpha, \beta, N) = y_s R_n^{(-1)}(y_s; \alpha, \beta, N),$$

where the grid y_s has the following expression

$$(3.3) \quad y_s = \begin{cases} -\alpha - \beta + 2s + 1 & \text{if } s \text{ even,} \\ \alpha + \beta - 2s - 1 & \text{if } s \text{ odd} \end{cases}$$

or, equivalently,

$$(3.4) \quad y_s = (-1)^s (1 - \alpha - \beta + 2s), \quad s = 0, 1, \dots, N.$$

The recurrence coefficients are

$$(3.5) \quad A_n^{(-1)} = \begin{cases} 2(n-N) & \text{if } n \text{ even} \\ 2(n+1-\alpha) & \text{if } n \text{ odd} \end{cases}, \quad C_n^{(-1)} = \begin{cases} -2n & \text{if } n \text{ even} \\ 2(N+1-\beta-n) & \text{if } n \text{ odd} \end{cases}.$$

The corresponding monic -1 dual Hahn polynomials satisfy relation (2.5), where

$$(3.6) \quad u_n^{(-1)} = A_{n-1}^{(-1)} C_n^{(-1)} = \begin{cases} 4n(\alpha-n) & \text{if } n \text{ even} \\ 4(N-n+1)(n+\beta-N-1) & \text{if } n \text{ odd} \end{cases}$$

and

$$(3.7) \quad b_n^{(-1)} = 1 - \alpha - \beta - A_n^{(-1)} - C_n^{(-1)} = \begin{cases} 2N+1-\alpha-\beta & \text{if } n \text{ even} \\ -2N-3+\alpha+\beta & \text{if } n \text{ odd} \end{cases}$$

It is convenient to introduce the " μ -numbers"

$$(3.8) \quad [n]_\mu = n + \mu(1 - (-1)^n),$$

which appear naturally in problems connected with the Dunkl operators [8]. One can then present the recurrence coefficients in the compact form

$$(3.9) \quad u_n^{(-1)} = 4[n]_\xi [N-n+1]_\eta, \quad b_n^{(-1)} = 2([n]_\xi + [N-n]_\eta) + 1 - \alpha - \beta,$$

where

$$\xi = \frac{\beta - N - 1}{2}, \quad \eta = \frac{\alpha - N - 1}{2}.$$

It is seen that $u_0 = u_{N+1} = 0$ as required for finite orthogonal polynomials. The positivity condition $u_n > 0, n = 1, 2, \dots, N$ is equivalent to the conditions

$$(3.10) \quad \alpha > N, \quad \beta > N.$$

These polynomials are orthogonal on the finite set of points y_s

$$(3.11) \quad \sum_{s=0}^N w_s R_n^{(-1)}(y_s) R_m^{(-1)}(y_s) = \kappa_0 u_1^{(-1)} u_2^{(-1)} \dots u_n^{(-1)} \delta_{nm},$$

where the discrete weights are defined as

$$(3.12) \quad w_{2s} = (-1)^s \frac{(-N/2)_s}{s!} \frac{(1-\alpha/2)_s (1-\alpha/2-\beta/2)_s}{(1-\beta/2)_s (N/2+1-\alpha/2-\beta/2)_s}, \quad s = 0, 1, 2, \dots, \frac{N}{2}$$

and

$$(3.13) \quad w_{2s+1} = (-1)^s \frac{(-N/2)_{s+1}}{s!} \frac{(1-\alpha/2)_s (1-\alpha/2-\beta/2)_s}{(1-\beta/2)_s (N/2+1-\alpha/2-\beta/2)_{s+1}}, \quad s = 0, 1, \dots, \frac{N}{2}-1.$$

The normalization coefficient is

$$(3.14) \quad \kappa_0 = \frac{\left(1 - \frac{\alpha+\beta}{2}\right)_{N/2}}{\left(1 - \frac{\beta}{2}\right)_{N/2}}.$$

Assume that $\alpha = N + \epsilon_1$, $\beta = N + \epsilon_2$, where $\epsilon_{1,2}$ are arbitrary positive parameters. This parametrization corresponds to the positive condition for the dual -1 Hahn polynomials. Then it is easily verified that all the weights are positive $w_s > 0$, $s = 0, 1, \dots, N$.

Moreover, the spectral points y_s are divided into two non-overlapping discrete sets of the real line:

$$\{1 - \delta, -3 - \delta, -7 - \delta, \dots, -2N + 1 - \delta\}$$

and

$$\{1 + \delta, 5 + \delta, 9 + \delta, \dots, 2N - 3 + \delta\},$$

where $\delta = \epsilon_1 + \epsilon_2 > 0$. The first set corresponds to y_s with even s and contains $1 + N/2$ points; the second set corresponds to y_s with odd s and contains $N/2$ points.

(ii) when $N = 1, 3, 5, \dots$ is odd, a nontrivial $q \rightarrow -1$ limit also exists iff $a \rightarrow -1$, $b \rightarrow -1$. We take the parametrization

$$(3.15) \quad q = -e^\varepsilon, \quad a = -e^{\alpha\varepsilon}, \quad b = -e^{\beta\varepsilon}, \quad \varepsilon \rightarrow 0$$

with real parameters α, β . Dividing again the recurrence relation (2.3) by $q + 1$ and taking the limit $\varepsilon \rightarrow 0$, we obtain the recurrence relation (3.2) where the grid y_s is defined as

$$(3.16) \quad y_s = \begin{cases} \alpha + \beta + 2s + 1 & \text{if } s \text{ even} \\ -\alpha - \beta - 2s - 1 & \text{if } s \text{ odd} \end{cases}$$

and the recurrence coefficients given by

$$(3.17) \quad A_n^{(-1)} = \begin{cases} 2(\alpha + n + 1) & \text{if } n \text{ even} \\ 2(n - N) & \text{if } n \text{ odd} \end{cases}, \quad C_n^{(-1)} = \begin{cases} -2n & \text{if } n \text{ even} \\ 2(\beta + N - n + 1) & \text{if } n \text{ odd} \end{cases}.$$

The corresponding monic -1 dual Hahn polynomials satisfy the standard relation (2.5), where

$$(3.18) \quad u_n^{(-1)} = A_{n-1}^{(-1)} C_n^{(-1)} = \begin{cases} 4n(N + 1 - n) & \text{if } n \text{ even} \\ 4(\alpha + n)(\beta + N + 1 - n) & \text{if } n \text{ odd} \end{cases}$$

and

$$(3.19) \quad b_n^{(-1)} = 1 + \alpha + \beta - A_n^{(-1)} - C_n^{(-1)} = \begin{cases} -1 - \alpha + \beta & \text{if } n \text{ even} \\ -1 + \alpha - \beta & \text{if } n \text{ odd} \end{cases}.$$

Again, as in the case of even N , it is possible to present the recurrence coefficients in the compact form

$$(3.20) \quad u_n^{(-1)} = 4[n]_\xi[N - n + 1]_\eta, \quad b_n^{(-1)} = 2([n]_\xi + [N - n]_\eta) - 2N - 1 - \alpha - \beta,$$

with $\xi = \alpha/2$, $\eta = \beta/2$. It is seen that in both cases: N even and N odd, the recurrence coefficients of the dual -1 Hahn polynomials are presented in the unified form (3.9) or (3.20) with the difference only residing with the parameters ξ, η .

It is seen that $u_0 = u_{N+1} = 0$ as required for finite orthogonal polynomials. The positivity condition $u_n > 0$, $n = 1, 2, \dots, N$ is equivalent either to condition

$$(3.21) \quad \alpha > -1, \beta > -1$$

or to condition $\alpha < -N$, $\beta < -N$. In what follows we shall use only condition (3.21).

The polynomials $R_n^{(-1)}(x)$ are orthogonal on the finite set of points y_s

$$(3.22) \quad \sum_{s=0}^N w_s R_n^{(-1)}(y_s) R_m^{(-1)}(y_s) = \kappa_0 u_1^{(-1)} u_2^{(-1)} \dots u_n^{(-1)} \delta_{nm},$$

where the discrete weights are defined as

$$(3.23) \quad w_{2s} = (-1)^s \frac{(-(N-1)/2)_s}{s!} \frac{(1/2 + \alpha/2)_s (1 + \alpha/2 + \beta/2)_s}{(1/2 + \beta/2)_s (N/2 + 3/2 + \alpha/2 + \beta/2)_s}, \quad s = 0, 1, 2, \dots, \frac{N-1}{2}$$

and

$$(3.24) \quad w_{2s+1} = (-1)^s \frac{(-(N-1)/2)_s}{s!} \frac{(1/2 + \alpha/2)_{s+1} (1 + \alpha/2 + \beta/2)_s}{(1/2 + \beta/2)_{s+1} (N/2 + 3/2 + \alpha/2 + \beta/2)_s}, \quad s = 0, 1, 2, \dots, \frac{N-1}{2}$$

The normalization coefficient is

$$(3.25) \quad \kappa_0 = \frac{\left(1 + \frac{\alpha+\beta}{2}\right)_{(N+1)/2}}{\left(\frac{\beta+1}{2}\right)_{(N+1)/2}}.$$

Assume that $\alpha = -1 + \epsilon_1$, $\beta = -1 + \epsilon_2$, where $\epsilon_{1,2}$ are arbitrary positive parameters. This parametrization corresponds to the positive condition for the dual -1 Hahn polynomials for N odd. Then it is easily verified that the weights are positive $w_s > 0$, $s = 0, 1, \dots, N$.

Moreover, the spectral points y_s are divided into two non-overlapped discrete sets of the real line:

$$\{-1 - \delta, -5 - \delta, -9 - \delta, \dots, -2N + 1 - \delta\}$$

and

$$\{-1 + \delta, 3 + \delta, 7 + \delta, \dots, 2N - 3 + \delta\},$$

where $\delta = \epsilon_1 + \epsilon_2 > 0$. Both sets contain $(N-1)/2$ points.

4. EXPLICIT EXPRESSION IN TERMS OF THE ORDINARY DUAL HAHN POLYNOMIALS

In this section, we derive an explicit expression for the -1 dual Hahn polynomials in terms of the ordinary dual Hahn polynomials.

Consider first the case of even $N = 2, 4, 6, \dots$. Introduce the "shifted" monic -1 dual Hahn polynomials $\tilde{R}_n(x) = R_n^{(-1)}(x-1)$. From formulas (3.6), (3.7), we conclude that these polynomials satisfy the recurrence relation

$$(4.1) \quad \tilde{R}_{n+1}(x) + (-1)^n \tau \tilde{R}_n(x) + u_n \tilde{R}_{n-1}(x) = x \tilde{R}_n(x),$$

where

$$\tau = 2N + 2 - \alpha - \beta$$

and u_n are given by (3.6).

A recurrence relation of the type (4.1) leads to orthogonal polynomials $\tilde{R}_n(x)$ which are very close to symmetric orthogonal polynomials. Using methods developed in [2] and [13], we can introduce a pair of monic orthogonal polynomials $P_n(x)$ and $Q_n(x)$ by the formulas :

$$(4.2) \quad \tilde{R}_{2n}(x) = P_n(x^2), \quad \tilde{R}_{2n+1}(x) = (x - \tau)Q_n(x^2).$$

It can easily be shown that the polynomials $P_n(x)$ and $Q_n(x)$ satisfy the following recurrence relations (it is assumed that $u_0 = 0$)

$$(4.3) \quad P_{n+1}(x) + (u_{2n} + u_{2n+1} + \tau^2)P_n(x) + u_{2n}u_{2n-1}P_{n-1}(x) = xP_n(x)$$

and

$$(4.4) \quad Q_{n+1}(x) + (u_{2n+2} + u_{2n+1} + \tau^2)Q_n(x) + u_{2n}u_{2n+1}Q_{n-1}(x) = xQ_n(x),$$

and moreover that the polynomials are connected by the Christoffel transform

$$(4.5) \quad Q_n(x) = \frac{P_{n+1}(x) + u_{2n+1}P_n(x)}{x - \tau^2}.$$

It is also easily seen that both $P_n(x)$ and $Q_n(x)$ are ordinary dual Hahn polynomials. We hence have the following explicit expression:

$$(4.6) \quad R_{2n}^{(-1)}(x-1) = \gamma_n^{(0)} {}_3F_2 \left(\begin{matrix} -n, \eta + \frac{x}{4}, \eta - \frac{x}{4} \\ -\frac{N}{2}, 1 - \frac{\alpha}{2} \end{matrix} \middle| 1 \right), \quad n = 0, 1, 2, \dots$$

and

$$(4.7) \quad R_{2n+1}^{(-1)}(x-1) = \gamma_n^{(1)} (x - \tau) {}_3F_2 \left(\begin{matrix} -n, \eta + \frac{x}{4}, \eta - \frac{x}{4} \\ 1 - \frac{N}{2}, 1 - \frac{\alpha}{2} \end{matrix} \middle| 1 \right), \quad n = 0, 1, 2, \dots,$$

where $\eta = 1/2 - (\alpha + \beta)/4$ and the normalization coefficients are

$$\gamma_n^{(0)} = 16^n (-N/2)_n (1 - \alpha/2)_n, \quad \gamma_n^{(1)} = 16^n (1 - N/2)_n (1 - \alpha/2)_n.$$

Quite similarly, for the odd $N = 1, 3, 5, \dots$ we find

$$(4.8) \quad R_{2n}^{(-1)}(x-1) = \gamma_n^{(0)} {}_3F_2 \left(\begin{matrix} -n, \eta + \frac{x}{4}, \eta - \frac{x}{4} \\ -\frac{N-1}{2}, \frac{\alpha+1}{2} \end{matrix} \middle| 1 \right), \quad n = 0, 1, 2, \dots$$

and

$$(4.9) \quad R_{2n+1}^{(-1)}(x-1) = \gamma_n^{(1)} (x + \alpha - \beta) {}_3F_2 \left(\begin{matrix} -n, \eta + \frac{x}{4}, \eta - \frac{x}{4} \\ -\frac{N-1}{2}, \frac{\alpha+3}{2} \end{matrix} \middle| 1 \right), \quad n = 0, 1, 2, \dots,$$

where

$$\eta = \frac{\alpha + \beta + 2}{4}, \quad \gamma_n^{(0)} = 16^n \left(\frac{1-N}{2} \right)_n \left(\frac{\alpha+1}{2} \right)_n, \quad \gamma_n^{(1)} = 16^n \left(\frac{1-N}{2} \right)_n \left(\frac{\alpha+3}{2} \right)_n.$$

Some of these polynomials have appeared in [9], [4] in the context of quantum spin chains.

5. DIFFERENCE EQUATION

Consider the following operator L defined on the space of functions $f(s)$ that depend on a discrete variable s :

$$(5.1) \quad Lf(s) = B(s)f(s+1) + D(s)f(s-1) - (B(s) + D(s))f(s), \quad s = 0, 1, 2, \dots,$$

where $B(s), D(s)$ are given in (2.7). Manifestly, the difference equation (2.6) means that the dual q -Hahn polynomials are eigenfunctions of the operator L

$$(5.2) \quad LR_n(x(s)) = \lambda_n R_n(x(s))$$

with the eigenvalues

$$(5.3) \quad \lambda_n = q^{-n} - 1.$$

When $q \rightarrow 1$ we obtain the difference eigenvalue equation for the ordinary dual Hahn polynomials

$$(5.4) \quad L_1 W_n(x(s)) = -n W_n(x(s)),$$

where the operator L_1 can be obtained from L as $L_1 = \lim_{q \rightarrow 1} L(q-1)^{-1}$. Explicitly [5]

$$(5.5) \quad L_1 = B_1(s)f(s+1) + D_1(s)f(s-1) - (B_1(s) + D_1(s))f(s),$$

where

$$B_1(s) = \frac{(s + \alpha + \beta + 1)(s + \alpha + 1)(N - s)}{(2s + \alpha + \beta + 1)(2s + \alpha + \beta + 2)}, \quad D_1(s) = \frac{s(s + \beta)(s + \alpha + \beta + N + 1)}{(2s + \alpha + \beta + 1)(2s + \alpha + \beta)}.$$

When we try to perform a similar procedure for the limit $q \rightarrow -1$ we encounter a problem. Indeed, it is easily seen that $L(1+q)^{-1}$ does not have a nondegenerate limit as $q \rightarrow -1$. We thus cannot obtain an eigenvalue equation in 3-diagonal form like (5.4) for the dual -1 Hahn polynomials.

Nevertheless we observe that the operator $(L^2 + 2L)(1+q)^{-1}$ does survive in the limit $q \rightarrow -1$. We hence have the following eigenvalue equation for the dual -1 Hahn polynomials

$$(5.6) \quad HR_n^{(-1)}(y_s) = 2nR_n^{(-1)}(y_s),$$

where the grid y_s is defined by (3.3) or (3.16) and

$$H = \lim_{q \rightarrow -1} (L^2 + 2L)(1+q)^{-1}.$$

The operator H obviously is 5-diagonal, i.e.

$$(5.7) \quad Hf(s) = U_2(s)(f(s+2) - f(s)) + U_1(s)(f(s+1) - f(s)) + V_2(s)(f(s-2) - f(s)) + V_1(s)(f(s-1) - f(s))$$

The explicit expressions for the coefficients $U_i(s), V_i(s)$ depend on the parity of N . For even $N = 2, 4, 6, \dots$ they are:

$$(5.8) \quad U_2(s) = \begin{cases} -2 \frac{(\alpha-s-2)(\beta+\alpha-s-2)(N-s)}{(\alpha+\beta-2s-2)(\alpha+\beta-2s-4)} & \text{if } s \text{ even} \\ -2 \frac{(\alpha-s-1)(\beta+\alpha-s-1)(N-s-1)}{(\alpha+\beta-2s-2)(\alpha+\beta-2s-4)} & \text{if } s \text{ odd} \end{cases},$$

$$(5.9) \quad U_1(s) = \begin{cases} 2 \frac{(\beta+\alpha)(\alpha-\beta)(N-s)}{(\alpha+\beta-2s)(\alpha+\beta-2s-2)(\alpha+\beta-2s-4)} & \text{if } s \text{ even} \\ 4 \frac{(\alpha-s-1)(\beta+\alpha-s-1)(2N+2-\alpha-\beta)}{(\alpha+\beta-2s)(\alpha+\beta-2s-2)(\alpha+\beta-2s-4)} & \text{if } s \text{ odd} \end{cases},$$

$$(5.10) \quad V_2(s) = \begin{cases} 2 \frac{s(\beta-s)(-\alpha-\beta+N+s)}{(\alpha+\beta-2s+2)(\alpha+\beta-2s)} & \text{if } s \text{ even} \\ 2 \frac{(s-1)(\beta-s+1)(-\alpha-\beta+N+s+1)}{(\alpha+\beta-2s+2)(\alpha+\beta-2s)} & \text{if } s \text{ odd} \end{cases},$$

$$(5.11) \quad V_1(s) = \begin{cases} 4 \frac{s(\beta-s)(2N+2-\alpha-\beta)}{(\alpha+\beta-2s)(\alpha+\beta-2s-2)(\alpha+\beta-2s+2)} & \text{if } s \text{ even} \\ -2 \frac{(\beta+\alpha)(\alpha-\beta)(-\alpha-\beta+N+s+1)}{(\alpha+\beta-2s)(\alpha+\beta-2s-2)(\alpha+\beta-2s+2)} & \text{if } s \text{ odd} \end{cases};$$

and for odd $N = 1, 3, 5, \dots$

$$(5.12) \quad U_2(s) = \begin{cases} -2 \frac{(\alpha+\beta+s+2)(\alpha+s+1)(N-s-1)}{(\alpha+\beta+2s+2)(\alpha+\beta+2s+4)} & \text{if } s \text{ even} \\ -2 \frac{(\alpha+s+2)(\alpha+\beta+s+1)(N-s)}{(\alpha+\beta+2s+2)(\alpha+\beta+2s+4)} & \text{if } s \text{ odd} \end{cases},$$

$$(5.13) \quad U_1(s) = \begin{cases} -2 \frac{(\alpha+\beta)(\alpha+s+1)(\alpha+\beta+2N+2)}{(\alpha+\beta+2s)(\alpha+\beta+2s+2)(\alpha+\beta+2s+4)} & \text{if } s \text{ even} \\ -4 \frac{(\alpha-\beta)(N-s)(\alpha+\beta+s+1)}{(\alpha+\beta+2s)(\alpha+\beta+2s+2)(\alpha+\beta+2s+4)} & \text{if } s \text{ odd} \end{cases},$$

$$(5.14) \quad V_2(s) = \begin{cases} -2 \frac{s(\beta+s-1)(\alpha+\beta+N+s+1)}{(\alpha+\beta+2s-2)(\alpha+\beta+2s)} & \text{if } s \text{ even} \\ -2 \frac{(s-1)(\beta+s)(\alpha+\beta+N+s)}{(\alpha+\beta+2s-2)(\alpha+\beta+2s)} & \text{if } s \text{ odd} \end{cases},$$

$$(5.15) \quad V_1(s) = \begin{cases} -4 \frac{s(\alpha-\beta)(\alpha+\beta+N+s+1)}{(\alpha+\beta+2s)(\alpha+\beta+2s+2)(\alpha+\beta+2s-2)} & \text{if } s \text{ even} \\ -2 \frac{(\beta+s)(\alpha+\beta)(\alpha+\beta+2N+2)}{(\alpha+\beta+2s)(\alpha+\beta+2s+2)(\alpha+\beta+2s-2)} & \text{if } s \text{ odd} \end{cases}.$$

We thus have a difference equation for the dual -1 Hahn polynomials in the form

$$\begin{aligned} & U_2(s) \left(R_n^{(-1)}(y_{s+2}) - R_n^{(-1)}(y_s) \right) + U_1(s) \left(R_n^{(-1)}(y_{s+1}) - R_n^{(-1)}(y_s) \right) + \\ & V_2(s) \left(R_n^{(-1)}(y_{s-2}) - R_n^{(-1)}(y_s) \right) + V_1(s) \left(R_n^{(-1)}(y_{s-1}) - R_n^{(-1)}(y_s) \right) = 2n R_n^{(-1)}(y_s), \end{aligned}$$

where the coefficients $U_i(s), V_i(s)$ are provided in the above formulas.

6. ANOTHER FORM OF THE DIFFERENCE EQUATION

The difference equation for the dual -1 Hahn polynomials can be presented in a more compact form if one notices that the grid y_s satisfies the relations

$$(6.1) \quad y_{s\pm 1} = \begin{cases} -y_s \mp 2 & \text{if } s \text{ even} \\ -y_s \pm 2 & \text{if } s \text{ odd} \end{cases}.$$

This property implies that the difference equation can be written as

$$E_1(x) \left(R_n^{(-1)}(x+4) - R_n^{(-1)}(x) \right) + E_2(x) \left(R_n^{(-1)}(x-4) - R_n^{(-1)}(x) \right) +$$

$$G_1(x) \left(R_n^{(-1)}(-x-2) - R_n^{(-1)}(x) \right) + G_2(x) \left(R_n^{(-1)}(-x+2) - R_n^{(-1)}(x) \right) = 2nR_n^{(-1)}(x)$$

or, in operator form as

$$(6.2) \quad HR_n^{(-1)}(x) = 2nR_n^{(-1)}(x).$$

The operator H in (6.2) reads

$$(6.3) \quad H = E_1(x)T^4 + E_2(x)T^{-4} + G_1(x)T^2R + G_2(x)T^{-2}R - (E_1(x) + E_2(x) + G_1(x) + G_2(x))I,$$

where the operators T and T^{-1} are the standard shift operators: $T^j f(x) = f(x+j)$, $j = 0, \pm 1, \pm 2, \dots$, and R is the reflection operator $Rf(x) = f(-x)$. I denotes the identity operator.

The functions $E_i(x), G_i(x)$, $i = 1, 2$ are simple rational functions in x .

For even $N = 2, 4, \dots$ we have

$$E_1(x) = \frac{(x+3-\alpha+\beta)(x+3-\alpha-\beta)(x-1-2N+\alpha+\beta)}{4(x+1)(3+x)},$$

$$E_2(x) = -\frac{(\alpha-1+\beta+x)(\alpha-1-\beta+x)(x-1+2N-\alpha-\beta)}{4(x-1)(x-3)},$$

$$G_1(x) = \frac{(\alpha^2 - \beta^2)(x+\alpha+\beta-2N-1)}{(x^2-1)(x+3)},$$

$$G_2(x) = \frac{(2N+2-\alpha-\beta)((x+\alpha-1)^2 - \beta^2)}{(x^2-1)(x-3)}$$

For odd $N = 1, 3, 5, \dots$

$$E_1(x) = \frac{(x+\alpha+\beta+3)(x+\alpha-\beta+1)(x-\alpha-\beta-2N+1)}{4(x+1)(x+3)}$$

$$E_2(x) = -\frac{(x-\alpha-\beta-1)(x+\beta-\alpha-3)(x+\alpha+\beta+2N+1)}{4(x-1)(x-3)}$$

$$G_1(x) = \frac{(\alpha+\beta)(\alpha+\beta+2+2N)(x+1+\alpha-\beta)}{(1-x^2)(x+3)}$$

$$G_2(x) = \frac{(\alpha-\beta)(x-\alpha-\beta-1)(x+\alpha+\beta+2N+1)}{(1-x^2)(x-3)}$$

The form (6.2) of the difference equation is preferable because we here have the operator H acting directly on the argument of the polynomials. The operator H belongs to the class of Dunkl shift operators: it contains both simple shifts T^j and the reflection operator R . Moreover, the operator H preserves the space of polynomials: it transforms any polynomials of degree n into a polynomial of the same degree n . Operators of this kind were considered in the theory of Bannai-Ito polynomials [11]. However, in contrast to the Bannai-Ito situation, the operator H in the present case is of *second order*, while the Bannai-Ito polynomials are eigenfunctions of a Dunkl-shift operator of the first order [11]. Equivalently, this means that for generic values of the parameters α, β , the dual -1 Hahn polynomials are eigenvectors of a 5-diagonal matrix (while the Bannai-Ito polynomials are eigenvectors of a 3-diagonal matrix).

Acknowledgments

The authors would like to gratefully acknowledge the hospitality extended to LV and AZ by Kyoto University and to ST and LV by the Donetsk Institute for Physics and Technology in the course of this investigation. The research of LV is supported in part by a research grant from the Natural Sciences and Engineering Research Council (NSERC) of Canada.

REFERENCES

- [1] E. Bannai and T. Ito, *Algebraic Combinatorics I: Association Schemes*. 1984. Benjamin & Cummings, Mento Park, CA.
- [2] T. Chihara, *On kernel polynomials and related systems*, Boll. Unione Mat. Ital., 3 Ser., **19** (1964), 451–459.
- [3] T. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, NY, 1978.
- [4] E. I. Jafarov, N. I. Stoilova and J. Van der Jeugt, *Finite oscillator models: the Hahn oscillator*, ArXiv: 1101.5310.
- [5] R. Koekoek, P. Lesky, R. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their Q -analogues*, Springer-Verlag, 2010.
- [6] D. Leonard, *Orthogonal Polynomials, Duality and Association Schemes*, SIAM J. Math. Anal. **13** (1982) 656–663.
- [7] A.F. Nikiforov, S.K. Suslov, and V.B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable*, Springer, Berlin, 1991.
- [8] M. Rosenblum, *Generalized Hermite Polynomials and the Bose-like Oscillator Calculus*, in: Oper. Theory Adv. Appl., vol. **73**, Birkhauser, Basel, 1994, pp. 369–396. ArXiv:math/9307224.
- [9] N. I. Stoilova and J. Van der Jeugt, *An Exactly Solvable Spin Chain Related to Hahn Polynomials*, SIGMA **7** (2011), 03.
- [10] P. Terwilliger, *Two linear transformations each tridiagonal with respect to an eigenbasis of the other*. Linear Algebra Appl. **330** (2001) 149–203.
- [11] S. Tsujimoto, L. Vinet and A. Zhedanov, *Dunkl-shift operators and Bannai-Ito polynomials*, arXiv:1106.3512.
- [12] L. Vinet and A. Zhedanov, *A “missing” family of classical orthogonal polynomials*, J. Phys. A: Math. Theor. **44** (2011) 085201, arXiv:1011.1669v2.
- [13] L. Vinet and A. Zhedanov, *A limit $q = -1$ for big q -Jacobi polynomials*, Trans.Amer.Math.Soc., to appear, arXiv:1011.1429v3.
- [14] L. Vinet and A. Zhedanov, *A Bochner theorem for Dunkl polynomials*, SIGMA, **7** (2011), 020; arXiv:1011.1457v3.

DEPARTMENT OF APPLIED MATHEMATICS AND PHYSICS, GRADUATE SCHOOL OF INFORMATICS,
KYOTO UNIVERSITY, SAKYO-KU, KYOTO 606–8501, JAPAN

CENTRE DE RECHERCHES MATHÉMATIQUES, UNIVERSITÉ DE MONTRÉAL, P.O. BOX 6128, CENTRE-
VILLE STATION, MONTRÉAL (QUÉBEC), H3C 3J7, CANADA

DONETSK INSTITUTE FOR PHYSICS AND TECHNOLOGY,, 83114 DONETSK, UKRAINE,